

Superscillations and transplanckian frequencies

H. Rosu*

Instituto de Física de la Universidad de Guanajuato, Apdo Postal E-143, León, Gto, México

Summary. - Superscillations can explain the arbitrarily high frequencies' paradox in black hole radiation.

PACS 04.60+n - Quantum theory of gravitation

LAA number: gr-qc/9606070

If one traces, in the usual quasi-geometrical derivation, the origin of the thermal radiation from a solar mass sized black hole a second after the formation of the black hole, its origin lies in the incoming vacuum frequencies of the order e^{10^4} (units are not important), implying scales of orders by far larger than the mass of the universe itself. This is almost a quotation from Unruh's work and represents the paradox of transplanckian frequencies, which is one of the most important unsolved issues in black hole radiation, and in fact, might be an indication of a serious drawback of the employed formalism. It has been first discussed by Jacobson [1] and by Unruh [2] in a hydrodynamical model. The conclusion of Unruh after a quite detailed analysis of two-dimensional 'dumb' holes is that there is an adiabatical adjustment of high frequency phenomena to low frequency physics close to the horizon (no high-frequency quantum hair). Since short distance cutoffs (at the Planck scale) are not affecting Hawking's thermal radiation, the common trend in the literature is to get rid of the transplanckian reservoir, e.g., by finding physical models (borrowed from other fields of physics) for the cutoffs [3].

Another way to get out from the transplanckian absurdity is to think of a material support of those extremely high frequencies, and if possible, of a physical entity and phenomenon related to black hole physics. Let me spell now my answer of principles to solve the transplanckian paradox, which I associate to *superscillations*, a remarkable phenomenon put forth by Aharonov [4]. Recall the underlined second in the first paragraph. This would correspond to a band-limited thermal signal of 1 Hz. Apparently, it is not well-known that band-limited functions can oscillate for arbitrarily long intervals arbitrarily faster than the highest frequency they contain. These are the superscillations. One can obtain them through a special two-parameter integral representation of band-limited functions (BLFs), which, in general, changes a BLF to a corresponding two-parameter superscillating function (SOF). As intuitively presented by Berry, the special integral representation of BLFs contains a Gaussian-like factor, which, in the zero-width limit acts like a δ -function and projects out SOFs. These SOFs appear in a subtle way in the 'quantum time-translation machine' concept of Aharonov, Anandan, Popescu, and Vaidman [4], but the present note has its origin in my

repeated reading of Professor Berry's works [5]. Among the physical examples of superscillations given by Berry, that of special interest for my argument here is the two-dimensional case of evanescent waves which can be expressed as the singular limit of an angular superposition of real plane waves. The main point now is that the in-out formalism does not include the quantum hair, i.e., the evanescent waves at the horizon. Any evanescent hair has the property that in the direction perpendicular to the exponentially decaying one (thus along the horizon), oscillates faster than the free-space wavenumber, forming a *superscillating reservoir*. One can construct angular superpositions of 'plane' waves for a Gaussian wave packet in the black hole background and show readily, just following Berry, that its singular limit is a superscillatory function [6]. Also, enclosed vortex configurations are important for the energy-momentum transport by means of evanescent modes.

Moreover, superscillations may be related to a well-known problem in signal processing, that of sampling a function faster than the Nyquist rate (Berry is quoting Daubechies), and also to quantum billiards, which would be very interesting to study in the black hole context. Berry gave an explicit example of reproducing Beethoven's ninth symphony (a signal spreading over 4000 s) with a signal bandlimited to 1 Hz by means of superscillations. As found by Berry, to reproduce Beethoven's ninth symphony with such superscillations requires a signal $e^{10^{19}}$ times stronger than with conventional oscillations. A similar feature shows up for black holes. In the specific case of a solar mass black hole to reproduce a one-second thermal symphony would require a signal e^{10^4} stronger in the superscillating part of the evanescent hair at the horizon. In other words, the incredible powerful signal is not to be found in the incoming vacuum frequencies but can be accommodated by various evanescent quantum hairs that any black hole possesses at the horizon. Thus, my claim is that black hole hairs, a not very exciting topic recently (due to the *no hair conjecture*), may prove useful in solving the transplanckian paradox, if considered as superscillating reservoirs.

Acknowledgments: This work was partially supported by the CONACyT project 4868-E9406. I thank Professor M.V. Berry for sending me his remarkable pa-

pers on superoscillations.

* Electronic-mail: rosu@ifug.ugto.mx

- [1] JACOBSON T., *Phys. Rev. D*, **44** (1991) 1731; “Introduction to black hole microscopy”, hep-th/9510026, to appear in *Proc. of the first Mexican school on gravitation and mathematical physics*, eds. A. Macias, T. Matos, O. Obregón and H. Quevedo (1996) .
- [2] UNRUH W.G., *Phys. Rev. D*, **51** (1995) 2827; *Phys. Rev. Lett.*, **46** (1981) 1351; see also, BROUT R. et al., hep-th/9506124 and CASHIER A. et al., hep-th/9606106 .
- [3] JACOBSON T., *Phys. Rev. D*, **53** (1996) 7082 .
- [4] AHARONOV Y., ANANDAN J., POPESCU S., and VAIDMAN L., *Phys. Rev. Lett.*, **64** (1990) 2965; See also, VAIDMAN L., *Found. Phys.*, **21** (1991) 947; SUTER D., *Phys. Rev. A*, **51** (1995) 45; VAIDMAN L., *Phys. Rev. A*, **52** (1995) 4297 .
- [5] BERRY MICHAEL, “Faster than Fourier”, in *Fundamental Problems in Quantum Theory*, eds. J.A. Anandan and J. Safko (Singapore, World Scientific), to appear, and *J. Phys. A: Math. Gen.*, **27** (1994) L391 .
- [6] ROSU H., work in progress .